



<https://moodle.lmu.de> → Kurse suchen: 'Rechenmethoden'

## Sheet 11: Delta Function and Fourier Series

Posted: Mo 10.01.22 Central Tutorial: Th 13.01.22 Due: Th 20.01.22, 14:00

(b)[2](E/M/A) means: problem (b) counts 2 points and is easy/medium hard/advanced

Suggestions for central tutorial: example problems 1, 3(a), 4, 5.

Videos exist for example problems 4 (C6.2.1), 5 (C6.3.5).

### Example Problem 1: Integrals with $\delta$ function [3]

Points: (a)[0.5](E); (b)[0.5](E); (c)[0.5](M); (d)[1](M); (e)[0.1](E).

Calculate the following integrals (with  $a \in \mathbb{R}$ ):

$$(a) \quad I_1(a) = \int_{-\infty}^{\infty} dx \delta(x - \pi) \sin(ax)$$

$$(b) \quad I_2(a) = \int_{\mathbb{R}^3} dx^1 dx^2 dx^3 \delta(\mathbf{x} - \mathbf{y}) \|\mathbf{x}\|^2, \quad \text{with } \mathbf{y} = (a, 1, 2)^T$$

$$(c) \quad I_3(a) = \int_0^a dx \delta(x - \pi) \frac{1}{a + \cos^2(x/2)}$$

$$(d) \quad I_4(a) = \int_0^3 dx \delta(x^2 - 6x + 8) \sqrt{e^{ax}}$$

$$(e) \quad I_5(a) = \int_{\mathbb{R}^2} dx^1 dx^2 \delta(\mathbf{x} - a\mathbf{y}) \mathbf{x} \cdot \mathbf{y}, \quad \text{with } \mathbf{y} = (1, 3)^T. \quad \text{Remark: } \delta(\mathbf{x}) = \delta(x^1)\delta(x^2).$$

[Check your results:  $I_1(\frac{1}{2}) = 1$ ,  $I_2(1) = 6$ ,  $I_3(\pi) = \frac{1}{2\pi}$ ,  $I_4(\ln 2) = 1$ ,  $I_5(1) = 10$ .]

### Example Problem 2: Lorentz representation of the Dirac $\delta$ -function [4]

Points: [4](M).

Explain why in the limit  $\epsilon \rightarrow 0^+$ , the Lorentz peak function  $\delta^\epsilon(x)$  given below is a representation of the Dirac delta function  $\delta(x)$ . To this end, compute (i) the height, (ii) the width  $x_w$  (defined by  $\delta^\epsilon(x_w) = \frac{1}{2}\delta^\epsilon(0)$ ,  $x_w > 0$ ) and (iii) the area of the peak. How do these quantities behave for  $\epsilon \rightarrow 0^+$ ? Furthermore, calculate the functions (iv)  $\Theta^\epsilon(x) = \int_{-\infty}^x dx' \delta^\epsilon(x')$  and (v)  $\delta'^\epsilon(x) = \frac{d}{dx}\delta^\epsilon(x)$ . Sketch  $\Theta^\epsilon$ ,  $\epsilon\delta^\epsilon$  and  $\epsilon^2\delta'^\epsilon$  as functions of  $x/\epsilon$  in three separate sketches (one beneath the other, with aligned  $y$ -axes and the same scaling for the  $x/\epsilon$ -axes).

$$\text{Lorentz-Peak: } \delta^\epsilon(x) = \frac{\epsilon/\pi}{x^2 + \epsilon^2}.$$

*Hint:* When calculating the peak weight, use the substitution  $x = \epsilon \tan y$ .

*Remark:* Lorentzian functions are common in physics. Example: the energy spectrum of a discrete quantum state, which is weakly coupled to the environment, has the form of a Lorentzian function, the width of which is determined by the strength of the coupling to the environment. As the coupling strength approaches zero, we obtain a  $\delta$  peak.

### Example Problem 3: Series representation of hyperbolic functions [3]

Points: [3](E).

Compute the following series for  $y \in \mathbb{R}^+$ , by expressing each as a geometric series in  $\omega \equiv e^{-y}$ .

$$(a) \sum_{n=0}^{\infty} e^{-y(n+1/2)}, \quad (b) \sum_{n=0}^{\infty} (-1)^n e^{-y(n+1/2)}, \quad (c) \sum_{n \in \mathbb{Z}} e^{-y|n|}.$$

### Example Problem 4: Fourier series of the sawtooth function [2]

Points: [2](M).

Let  $f(x)$  be a sawtooth function, defined by  $f(x) = x$  for  $-\pi < x < \pi$ ,  $f(\pm\pi) = 0$  and  $f(x+2\pi) = f(x)$ . Calculate the Fourier coefficients  $\tilde{f}_n$  in the representation  $f(x) = \frac{1}{L} \sum_n e^{ik_n x} \tilde{f}_n$ . How should  $k_n$  and  $L$  be chosen? Sketch the function  $f(x)$ , as well as the sum of the  $n = 1$  and  $n = -1$  terms of the Fourier series (i.e. the first term of the corresponding sine series). [Check your result:  $\tilde{f}_6 = \frac{1}{3}i\pi$ .]

### Example Problem 5: Parseval's identity and convolution [7]

Points: (a)[3](M); (b)[2](M); (c)[2](M).

Let  $f(x)$  be a sawtooth function, defined by  $f(x) = x$  for  $-\pi < x < \pi$ ,  $f(\pm\pi) = 0$  and  $f(x+2\pi) = f(x)$ . In the Fourier representation  $f(x) = \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} e^{inx} \tilde{f}_n$ , its Fourier coefficients are  $\tilde{f}_0 = 0$ ,  $\tilde{f}_{n \neq 0} = 2\pi i (-1)^n / n$ . (See example problem 4.) Let  $g(x) = \sin x$ .

- Using this concrete example, check that Parseval's identity holds, by computing both the integral  $\int_{-\pi}^{\pi} dx \bar{f}(x)g(x)$  and the sum  $(1/2\pi) \sum_n \bar{\tilde{f}}_n \tilde{g}_n$  explicitly.
- Prove the famous identity  $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$ , by computing the integral  $\int_{-\pi}^{\pi} dx f^2(x)$  in two ways: first, by direct integration, and second, by expressing it as a sum over Fourier modes using Parseval's identity.
- Calculate the convolution  $(f * g)(x)$  both by directly computing the convolution integral and by using the convolution theorem and a summation of Fourier coefficients.

---

[Total Points for Example Problems: 19]

---

### Homework Problem 1: Integrals with $\delta$ function [4]

Points: (a)[0.5](E); (b)[0.5](E); (c)[0.5](M); (d)[1](M); (e)[1](A); (f)[0.5](E).

Calculate the following integrals (with  $a \in \mathbb{R}$ ,  $n \in \mathbb{N}$ ):

$$(a) \quad I_1(a) = \int_1^4 dx \delta(x-2) (a^x + 3)$$

$$(b) \quad I_2(a) = \int_{\mathbb{R}^2} dx^1 dx^2 \delta(\mathbf{x} - \mathbf{y}) (x^1 + x^2)^2 e^{3-x^1}, \quad \text{with } \mathbf{y} = (3, a)^T$$

$$(c) \quad I_3(a) = \int_{-1}^1 dx \sqrt{2+2x} \delta(ax-2), \quad \text{with } a \neq 0$$

$$(d) \quad I_4(a) = \int_{-\infty}^{\infty} dx \delta(3^{-x} - 9)(1 - x^a)$$

$$(e) \quad I_5(n) = \int_{-\pi/2}^{9\pi/2} dx \cos(nx) \delta(\sin x)$$

$$(f) \quad I_6(a) = \int_{\mathbb{R}^2} dx^1 dx^2 \delta(\mathbf{x} - \mathbf{y}) e^{\|\mathbf{x}\|^2}, \quad \text{with } \mathbf{y} = (a, -a)^T$$

[Check your results:  $I_1(3) = 12$ ,  $I_2(-5) = 4$ ,  $I_3(2) = \frac{1}{2}$ ,  $I_4(3) = \frac{1}{\ln 3}$ ,  $I_5(7) = 1$ ,  $I_6(\frac{1}{\sqrt{2}}) = e$ .]

### Homework Problem 2: Representations of the Dirac $\delta$ -function [4]

Points: [4](M).

Explain why in the limit  $\epsilon \rightarrow 0^+$ , the peak-shaped function  $\delta^\epsilon(x)$  given below is a representation of the Dirac delta function  $\delta(x)$ . To this end, compute (i) the height, (ii) the width  $x_w$  (defined by  $\delta^\epsilon(x_w) = \frac{1}{2}\delta^\epsilon(0)$ ,  $x_w > 0$ ) and (iii) the area of the peak. How do these quantities behave for  $\epsilon \rightarrow 0^+$ ? Furthermore, calculate the functions (iv)  $\Theta^\epsilon(x) = \int_{-\infty}^x dx' \delta^\epsilon(x')$  and (v)  $\delta'^\epsilon(x) = \frac{d}{dx} \delta^\epsilon(x)$ . Sketch  $\Theta^\epsilon$ ,  $\epsilon \delta^\epsilon$  and  $\epsilon^2 \delta'^\epsilon$  as functions of  $x/\epsilon$  in three separate sketches (one beneath the other, with aligned  $y$ -axes and the same scaling for the  $x/\epsilon$ -axes).

$$\text{Derivative of the Fermi function: } \delta^\epsilon(x) = \frac{1}{4\epsilon} \frac{1}{\cosh^2[x/(2\epsilon)]}.$$

*Hint:* When calculating the peak weight, use the substitution  $y = \tanh[x/(2\epsilon)]$ .

*Remark:* In condensed matter physics and nuclear physics the function  $\delta^\epsilon(x)$  plays an important role: it arises as the derivative of the so-called **Fermi function**,  $f(E) = \frac{1}{e^{E/k_B T} + 1} = \Theta^{k_B T}(-E)$ , with  $-\frac{d}{dE} f(E) = \delta^{k_B T}(E)$ , where  $f(E)$  is the occupation probability of a fermionic single-particle state with energy  $E$  as function of the system's temperature  $T$  ( $k_B$  is the so-called Boltzmann constant). In the limit of zero temperature,  $T \rightarrow 0$ , the derivative of the Fermi function reduces to a Dirac  $\delta$ -function.

### Homework Problem 3: Series representation of the periodic $\delta$ function [5]

Points: (a)[0.5](E); (b)[0.5](M); (c)[1.5](A); (d)[0.5](E); (e)[1](A); (f)[0.5](E); (g)[0.5](E)

Show that the function  $\delta^\epsilon(x)$ , defined by

$$\delta^\epsilon(x) = \frac{1}{L} \sum_k e^{ikx - \epsilon|k|}, \quad k = 2\pi n/L, \quad n \in \mathbb{Z}, \quad x, \epsilon, L \in \mathbb{R}, \quad 0 < \epsilon \ll L, \quad (1)$$

has the following properties:

$$(a) \quad \delta^\epsilon(x) = \delta^\epsilon(x + L). \quad (2)$$

$$(b) \quad \int_{-L/2}^{L/2} dx \delta^\epsilon(x) = 1. \quad \text{Hint: Treat } k = 0 \text{ and } k \neq 0 \text{ separately in } \sum_k. \quad (3)$$

$$(c) \quad \delta^\epsilon(x) = \frac{1}{2L} \left[ \frac{1+w}{1-w} + \frac{1+\bar{w}}{1-\bar{w}} \right] = \frac{1}{L} \frac{1 - e^{-4\pi\epsilon/L}}{1 + e^{-4\pi\epsilon/L} - 2e^{-2\pi\epsilon/L} \cos(2\pi x/L)}, \quad (4)$$

where  $w = e^{2\pi(ix-\epsilon)/L}$  and  $\bar{w} = e^{2\pi(-ix-\epsilon)/L}$ .

Hint: Write out the sum in Eq. (1) as a geometric series in powers of  $w$  and  $\bar{w}$ .

$$(d) \quad \lim_{\epsilon \rightarrow 0} \delta^\epsilon(x) = 0 \quad \text{for } x \neq mL, \text{ with } m \in \mathbb{Z}. \quad \text{Hint: Start from Eq. (4).}$$

$$(e) \quad \delta^\epsilon(x) \simeq \frac{\epsilon/\pi}{\epsilon^2 + x^2} \quad \text{for } |x|/L \ll 1 \text{ and } \epsilon/L \ll 1.$$

Hint: Taylor expand the numerator in Eq. (4) up to first order in  $\tilde{\epsilon} = 2\pi\epsilon/L$ , and the denominator up to second order in  $\tilde{\epsilon}$  and  $\tilde{x} = 2\pi x/L$ .

(f) Sketch the function  $\delta^\epsilon(x)$  qualitatively for  $\epsilon/L \ll 1$  and  $x \in [-\frac{7}{2}L, \frac{7}{2}L]$ .

(g) Deduce that in the limit of  $\epsilon \rightarrow 0$ ,  $\delta^\epsilon(x)$  represents a periodic  $\delta$  function, with

$$\delta^0(x) = \frac{1}{L} \sum_k e^{ikx} = \sum_{m \in \mathbb{Z}} \delta(x - mL).$$

#### Homework Problem 4: Fourier series [4]

Points: (a)[2](E); (b)[2](M)

Determine the Fourier series for the following periodic functions, i.e. calculate the Fourier coefficients  $\tilde{f}_n$  in the representation  $f(x) = \frac{1}{L} \sum_n e^{ik_n x} \tilde{f}_n$ . How should  $k_n$  and  $L$  be chosen in each case? Sketch the functions first.

$$(a) \quad f(x) = |\sin x|, \quad (b) \quad f(x) = \begin{cases} 4x & \text{for } -\pi \leq x < 0, \\ 2x & \text{for } 0 \leq x < \pi, \end{cases} \quad \text{and } f(x + 2\pi) = f(x).$$

[Check your results: (a)  $\tilde{f}_3 = -\frac{2}{35}$ , (b)  $\tilde{f}_3 = \frac{2}{9}(2 - 9i\pi)$ .]

#### Homework Problem 5: Computing an infinite series using the convolution theorem [1]

Points: (a)[0.5](E); (b)[0.5](M); (c)[2](A,Bonus)

This problem illustrates how a complicated sum may be calculated explicitly using the convolution theorem.

Consider the periodic function  $f_\gamma(t) = f_\gamma(0)e^{\gamma t}$  for  $t \in [0, \tau)$  and  $f(t + \tau) = f(t)$ , with  $f_\gamma(0) = 1/(e^{\gamma\tau} - 1)$ . Take both  $\gamma$  and  $\tau$  to be positive numbers, so that  $f_{\pm\gamma}(0) \gtrless 0$ .

(a) Consider a Fourier series representation of  $f_\gamma(t)$  of the following form:

$$f_\gamma(t) = \frac{1}{\tau} \sum_{\omega_n} e^{-i\omega_n t} \tilde{f}_{\gamma,n}, \quad \tilde{f}_{\gamma,n} = \int_0^\tau dt e^{i\omega_n t} f_\gamma(t), \quad \text{with } \omega_n = 2\pi n/\tau, \quad n \in \mathbb{Z}.$$

Show that the Fourier coefficients are given by  $\tilde{f}_{\gamma,n} = 1/(i\omega_n + \gamma)$ .

- (b) Use this result and the convolution theorem to express the following series as a convolution of  $f_\gamma$  and  $f_{-\gamma}$ :

$$S(t) = \sum_{n=-\infty}^{\infty} \frac{e^{-i\omega_n t}}{\omega_n^2 + \gamma^2} = -\tau \int_0^\tau dt' f_\gamma(t-t') f_{-\gamma}(t'). \quad (5)$$

- (c) Sketch the functions  $f_\gamma(t-t')$  and  $f_{-\gamma}(t')$  occurring in the convolution theorem as functions of  $t'$ , for  $t' \in [-\tau, 2\tau]$ . Assume  $0 \leq t \leq \tau$  and show that the convolution integral (5) is given by the following expression:

$$S(t) = \frac{\tau [\sinh(\gamma(t-\tau)) - \sinh(\gamma t)]}{2\gamma [1 - \cosh(\gamma\tau)]}.$$

*Hint:* The integral  $\int_0^\tau dt'$  involves an interval of  $t'$  values for which  $t-t'$  lies outside of  $[0, \tau]$ . It is therefore advisable to split the integral into two parts, with  $\int_0^t dt'$  and  $\int_t^\tau dt'$ .

---

[Total Points for Homework Problems: 18]

---